arXiv:1010.3934v1 [math.AP] 19 Oct 2010

MULTI-ANISOTROPIC GEVREY REGULARITY OF HYPOELLIPTIC OPERATORS

CHIKH BOUZAR AND AHMED DALI

Abstract. We show a multi-anisotropic Gevrey regularity of solutions of hypoelliptic equations. This result is a precision of a classical result of Hörmander

1. Introduction

An important problem among others of linear partial differential equations is the $\mathcal{C}^{\infty}$ or Gevrey regularity of solutions of these equations. L. Hörmander has completely characterized the $\mathcal{C}^{\infty}$ regularity (hypoellipticity) of linear partial differential operators with complex constant coefficients, see [7]. An another fundamental result obtained by L. Hörmander says that every hypoelliptic differential operator $P(D)$ is anisotropic Gevrey hypoelliptic, i.e. $\exists \varrho = (\varrho_1, ..., \varrho_n) \in \mathbb{R}^n_+$ such that

$$u \in \mathcal{D}'(\Omega) \ \ and \ P(D)u = 0 \Longrightarrow \ u \in G^{\varrho}(\Omega),$$

where $G^{\varrho}(\Omega)$ is an anisotropic Gevrey space associated with $P(D)$.

A large class of hypoelliptic differential operators is the class of multi-quasielliptic differential operators, see V. P. Mikhaïlov [8], J. Friberg [3] and S. G. Gindikin, L. R. Volevich [4].

L. Zanghirati [10], proved that multi-quasielliptic differential operators are multi-anisotropic Gevrey hypoelliptic, i.e.

$$u \in \mathcal{D}'(\Omega) \ \ and \ P(D)u \in G^{s,\,\Gamma}(\Omega) \ \ \Longrightarrow \ u \in G^{s,\,\Gamma}(\Omega),$$

where $G^{s,\,\Gamma}(\Omega)$ is a Gevrey multi-anisotropic space associated with $P(D)$. This result clarifies the classical result of Hörmander in the case of multi-quasielliptic operators. The result of L. Zanghirati has been extended by C. Bouzar and R. Chaïli in [1] to multi-quasielliptic systems of differential operators.

The aim of this paper is to prove the multi-anisotropic Gevrey regularity of hypoelliptic linear differential operators with complex constant coefficients, and consequently we precise the result of Hörmander and extend the result of Zanghirati.

2. Multi-quasiellipticity

Let Ω be an open subset of $\mathbb{R}^n$, if $\alpha = (\alpha_1, ...\alpha_n) \in \mathbb{Z}^n_+$, $q = (q_1, .., q_n) \in \mathbb{R}^n_+$ and $\xi = (\xi_1, ..., \xi_n)$ $\in \mathbb{R}^n$, we set

$$|\alpha| = \alpha_1 + ... + \alpha_n$$

$$< \alpha, q >= \sum_{j=1}^{n} \alpha_j q_j$$

1991 *Mathematics Subject Classification.* Primary 35H10; Secondary 35D10, 35H30.

Key words and phrases. Hypoelliptic operators, Gevrey regularity, Multi-anisotropic Gevrey spaces, Newton polyhedron, Multi-quasiellipticity, Gevrey vectors .

$$\xi^{\alpha} = \xi_1^{\alpha_1} ... \ \xi_n^{\alpha_n}$$

$$D^{\alpha} = D_1^{\alpha_1} ... \ D_n^{\alpha_n}, D_j = \frac{1}{i}\frac{\partial}{\partial \xi_j}, \ j = 1, ..., n.$$

$$\mathbb{R}_+^n = \{\xi \in \mathbb{R}^n : \xi_j > 0, \ j = 1, .., n\}$$

The space $\mathcal{C}_0^{\infty}(\Omega)$ is the space of functions $u \in \mathcal{C}^{\infty}$ with compact support in Ω. The space of distributions on Ω is denoted $\mathcal{D}'(\Omega)$.

Definition 1. *Let A be a finite subset of $\overline{\mathbb{R}_+^n}$, the Newton's polyhedron of A, denoted $\Gamma(A)$, is the convex hull of $\{0\} \cup A$.*

A Newton's polyhedron Γ is always characterized by

$$\Gamma = \bigcap_{q \in \mathcal{A}(\Gamma)} \left\{\alpha \in \overline{\mathbb{R}_+^n}, \ \langle q, \alpha \rangle \leq 1\right\},$$

where $\mathcal{A}(\Gamma)$ is a finite subset of $\mathbb{R}^n \ \{0\}$.

Definition 2. *Let $\Gamma = \bigcap\limits_{q \in \mathcal{A}(\Gamma)} \left\{\alpha \in \overline{\mathbb{R}_+^n}, \ \langle q, \alpha \rangle \leq 1\right\}$ be a Newton's polyhedron, Γ is said to be regular, if*

$$q_j > 0, \ \forall j = 1, ..., n; \ \forall q = (q_1, ..., q_n) \in \mathcal{A}(\Gamma)$$

We associate with a regular Newton's polyhedron Γ the following elements

$$\mathcal{V}(\Gamma) = \left\{s^0 = 0, \ s^1, ..., s^{m(\Gamma)}\right\} \text{ the set of vertices of } \Gamma$$

$$|\xi|_{\Gamma} = \sum_{\nu \in \mathcal{V}(\Gamma)} |\xi|^{\nu}, \ \xi \in \mathbb{R}^n, \text{ where } |\xi|^{\nu} = |\xi_1|^{\nu_1} ... |\xi_n|^{\nu_n}$$

$$k(\alpha, \Gamma) = \inf\left\{t > 0, \ t^{-1}\alpha \in \Gamma\right\} = \max_{q \in \mathcal{A}(\Gamma)} \langle \alpha, q \rangle, \ \alpha \in \mathbb{R}_+^n$$

$$\mu(\Gamma) = \max_{\substack{q \in \mathcal{A}(\Gamma) \\ 1 \leq j \leq n}} q_j^{-1} \text{ called the formal order of } \Gamma$$

A differential operators with complex constant coefficients

$$P(D) = \sum_{\alpha} a_{\alpha} D^{\alpha}$$

has its complete symbol

$$P(\xi) = \sum_{\alpha} a_{\alpha} \xi^{\alpha}$$

Definition 3. *The Newton's polyhedron of P, denoted $\Gamma(P)$, is the convex hull of the set $\{0\} \cup \left\{\alpha \in \mathbb{Z}_+^n : \ a_{\alpha} \neq 0\right\}$.*

Define the weight function

$$|\xi|_{\mathbb{P}} = \sum_{\alpha \in \mathcal{V}(P)} |\xi^{\alpha}|, \ \forall \xi \in \mathbb{R}^n,$$

where $\mathcal{V}(P) = \mathcal{V}(\Gamma(P))$ is the set of vertices of $\Gamma(P)$. Recall

$$d(\xi) := dist(\xi, N(P)), \text{ where } N(P) := \{\zeta \in \mathbb{C}^n : P(\zeta) = 0\}$$

Definition 4. *The differential operator $P(D)$ is said hypoelliptic in Ω, if*

$$singsuppP(D)u = singsuppu,\ \forall u \in \mathcal{D}'(\Omega)$$

The characterization of hypoelliptic differential operators with constant coefficients is du to L. Hörmander. The following result, see the theorem 4.1.3 of [7], gives some characterizations of the hypoellipticity.

Theorem 1. *Let $P(D)$ be a differential operator with constant coefficients, the following properties are equivalent :*

i) The operator $P(D)$ is hypoelliptic.

ii) $\exists C > 0,\ \exists d > 0,\ |\xi|^d \leq Cd(\xi),\ \forall \xi \in \mathbb{R}^n,\ |\xi|$ large.

iii) If $\xi \in \mathbb{R}^n, |\xi| \to +\infty$, then $\dfrac{|D^\alpha P(\xi)|}{|P(\xi)|} \to 0,\ \forall \alpha \neq 0.$

iv) $\exists C > 0,\ \exists \rho > 0,\ \dfrac{|D^\alpha P(\xi)|}{|P(\xi)|} \leq C\,|\xi|^{-\rho|\alpha|},\ \forall \xi \in \mathbb{R}^n,\ |\xi|$ large.

The connection between an hypoelliptic operator and its Newton's polyhedron is given by the following proposition.

Proposition 2. *The Newton's polyhedron of an hypoelliptic differential operator is regular.*

Proof. See [3]. □

Remark 1. *The converse is not true, $\square = D_x^2 - D_y^2$ has a regular Newton's polyhedron with vertices $\{(0,0), (2,0), (0,2)\}$, but the operator $\square$ is not hypoelliptic.*

We introduce multi-quasielliptic polynomials which are a natural generalization of the classical quasi-elliptic operators. These operators were characterized first by V. P. Mikhaïlov [8], then studied by J. Friberg [3] and finally far developed by S. G. Gindikin and L. R. Volevich [4].

Definition 5. *The polynomial $P(\xi) = \sum\limits_{\alpha} a_\alpha \xi^\alpha$ is said to be multi-quasielliptic, if*

i) *its Newton's polyhedron $\Gamma(P)$ is regular.*

ii) $\exists C > 0$ *such that*

$$|\xi|_{\mathbb{P}} \leq C(1 + |P(\xi)|),\ \forall \xi \in \mathbb{R}^n$$

Proposition 3. *A multi-quasielliptic operator $P(D)$ is hypoelliptic.*

Proof. See [3] or [4]. □

Remark 2. *The converse is not true. Indeed, consider the following polynomial*

$$\begin{aligned} P(\xi,\eta) &= i\xi^5 + i\xi\eta^4 - 4i\xi^4\eta - 4i\xi^2\eta^3 + 6i\xi^3\eta^2 + i\xi^3 + i\xi\eta^2 + \xi^4\eta^2 \\ &\quad + \eta^6 - 4\xi^3\eta^3 - 4\xi\eta^5 + 6\xi^2\eta^4 + \eta^2\xi^2 + \eta^4, \end{aligned}$$

which is hypoelliptic thanks to the theorem 4.1.9 of [7]. We have

$$P_{(1,1)}(\xi,\eta) = \eta^2\left(\xi^4 + \eta^4 - 4\xi^3\eta - 4\xi\eta^3 + 6\xi^2\eta^2\right) = \eta^2(\xi - \eta)^4$$

The $q = (1,1)-$quasiprincipal part of $P(\xi,\eta)$ degenerates on the straight $\xi = \eta$, hence the polynomial $P(\xi,\eta)$ is not multi-quasielliptic, see [4].

3. Multi-anisotropic Gevrey vectors

The multi-anisotropic Gevrey spaces were explicitly defined by L. Zanghirati in [10] for studying the multi-anistropic Gevrey regularity of multi-quasielliptic differential operators by the method of elliptic iterates.

Definition 6. *Let Ω be an open subset of $\mathbb{R}^n$, Γ a regular Newton's polyhedron and $s \geq 1$. Denote $G^{s,\,\Gamma}(\Omega)$ the space of functions $u \in \mathcal{C}^\infty(\Omega)$ such that $\forall K \subset \Omega, \exists C > 0$, $\forall \alpha \in \mathbb{Z}_+^n$,*

$$\sup_{x\in K} |D^\alpha u(x)| \leq C^{|\alpha|+1} k(\alpha,\Gamma)^{s\mu\, k(\alpha,\Gamma)} \tag{1}$$

Example 1. *If Γ is the regular Newton's polyhedron defined by*

$$\Gamma = \left\{ \alpha \in \overline{\mathbb{R}_+^n} : \sum_{j=1}^n m_j^{-1}\alpha_j \leq 1, m_j \in \mathbb{R}_+ \right\},$$

then

$$G^{s,\,\Gamma}(\Omega) = \left\{ \begin{array}{c} u \in \mathcal{C}^\infty(\Omega), \forall K \subset \Omega, \exists C > 0, \forall \alpha \in \mathbb{Z}_+^n \\ |D^\alpha u(x)| \leq C^{|\alpha|+1} \langle \alpha, q\rangle^{s\langle \alpha, q\rangle} \end{array} \right\},$$

where $q := \left(\dfrac{m}{m_1}, ..., \dfrac{m}{m_n}\right)$ and $m := \max_j m_j$, i.e. $G^{s,\,\Gamma}(\Omega)$ is the classical anisotropic Gevrey space $G^{s,\,q}(\Omega)$. If $m_1 = m_2 = ... = m_n$, we obtain the classical isotropic Gevrey space $G^s(\Omega)$.

Definition 7. *Let Γ be the regular Newton's polyhedron of $P(D)$ and $s \geq 1$, the space of Gevrey vectors of $P(D)$, denoted $G^s(\Omega, P)$, is the space of $u \in \mathcal{C}^\infty(\Omega)$ such that , $\forall K$ compact of Ω, $\exists C > 0, \forall l \in \mathbb{N}$,*

$$\left\| P^l u \right\|_{L^\infty(K)} \leq C^{l+1} (l!)^{s\mu(\Gamma)}$$

Remark 3. *We can take $l^{sl\mu(\Gamma)}$ instead of $(l!)^{s\mu(\Gamma)}$.*

We recall a result of L. Zanghirati [10] and C. Bouzar and R. Chaïli [1] wich gives the multi-anisotropic Gevrey regularity of Gevrey vectors of multi-quasielliptic operators.

Theorem 4. *Let Ω be an open subset of $\mathbb{R}^n$, $s > 1$ and P a linear differential operator with complex constant coefficients with regular Newton's polyhedron Γ. Then the following assertions are equivalent :*
i) P is multi-quasielliptic in Ω
ii) $G^s(\Omega, P) = G^{s,\,\Gamma}(\Omega)$

4. Multi-anisotropic Gevrey hypoellipticity of hypoelliptic operators

In this section, $P = \sum_\alpha a_\alpha D^\alpha$ is an hypoelliptic differential operator with complex constant coefficients.

Definition 8. *A finite set $\mathcal{H} \subset \overline{\mathbb{R}_+^n}$ is said a polyhedron of hypoellipticity of P, if*

1. *$\forall \nu \in \mathcal{H}, \exists C > 0, \forall \xi \in \mathbb{R}^n$, $|\xi|^\nu \leq C(1 + d(\xi))$*
2. *$\mathcal{H}$ has vertices with rational components.*
3. *$\mathcal{H}$ is regular.*

Remark 4. *If* ν *belongs to the convex hull of* $\mathcal{H}$, *i.e.* $\nu = \sum\limits_{\substack{i\in I \\ I\ fini}} \lambda_i \beta_i$, *where* $\beta_i \in \mathcal{H}$ *and* $\sum\limits_{i\in I} \lambda_i = 1, \lambda_i \geq 0$, *then* $|\xi|^{\nu} \leq C\left(1 + d(\xi)\right), \forall \xi \in \mathbb{R}^n$, *therefor it is natural to assume that* $\mathcal{H}$ *is convex.*

Remark 5. *The set* $\mathcal{H}$ *is never empty, as an hypoelliptic operator satisfies :* $\exists C > 0,\ \exists d > 0, |\xi|^{d} \leq C\left(1 + d\left|\xi\right|\right), \forall \xi \in \mathbb{R}^n$.

Definition 9. *Denote* σ *be the smallest natural integer such that*

$$\sigma \mathcal{V}\left(\mathcal{H}\right) \subset 2\mathbb{N}_0^n,$$

and define the differential operator $Q_{\mathcal{H}}\left(D\right)$, *by*

$$Q_{\mathcal{H}}\left(D\right) = \sum_{\alpha \in \mathcal{V}(\mathcal{H})} D^{\sigma\alpha}$$

Proposition 5. *The operator* $Q_{\mathcal{H}}\left(D\right)$ *is multi-quasielliptic.*

Proof. The Newton's polyhedron of the differential operator $Q_{\mathcal{H}}$ has vertices with even positive integer components. Then

$$\left|Q_{\mathcal{H}}\left(\xi\right)\right| = \sum_{\alpha \in \mathcal{V}(\mathcal{H})} \left|\xi^{\sigma\alpha}\right| = \left|\xi\right|_{Q_{\mathcal{H}}},$$

hence

$$1 + \left|\xi\right|_{Q_{\mathcal{H}}} \leq \left(1 + \left|Q_{\mathcal{H}}\left(\xi\right)\right|\right),\ \forall \xi \in \mathbb{R}^n$$

□

Let $v \in \mathcal{C}_0^{\infty}\left(\mathbb{R}^n\right), s \in \mathbb{Z}_+$ and $\varepsilon > 0$, then

$$|||v|||_{s,\varepsilon}^2 := \int_{\mathbb{R}^n} \left(1 + \varepsilon d\left(\xi\right)\right)^s \left|\widehat{v}\left(\xi\right)\right|^2 d\xi$$

The following result is the lemma 4.4.3 of [7].

Lemma 6. *Let* u *be a solution of the equation* $Pu = 0$ *defined in the ball* $B_\varepsilon = \left\{x \in \mathbb{R}^n : |x| < \varepsilon\right\}$, *and let* $\varphi \in \mathcal{C}_0^{\infty}\left(B_1\right)$ *and the integer* $s \geq 1$. *Then*

$$\sum_{\alpha \neq 0} \varepsilon^{-2|\alpha|} \left|\left|\left|P^{(\alpha)}\left(D\right)\left(\varphi^{\varepsilon} u\right)\right|\right|\right|_{s,\varepsilon}^2 \leq C \sum_{\alpha \neq 0} \varepsilon^{-2|\alpha|} \int_{B_\varepsilon} \left|P^{(\alpha)}\left(D\right) u\right|^2 dx, \tag{2}$$

where C *is independent of* ε *and* u.

Remark 6. *In the lemma* φ^{ε} *denotes* $\varphi^{\varepsilon}\left(x\right) := \varphi\left(\frac{x}{\varepsilon}\right)$.

Thanks to this lemma, we obtain the following result.

Lemma 7. *Let* $\beta \in \mathbb{Z}_+^n \cap \sigma\mathcal{H}$, *then there exists a constant* $C > 0$, *such that for every solution* u *of* $Pu = 0$ *in* B_ε *and* $\varepsilon \in \left]0, 1\right[$, *we have*

$$\varepsilon^{2\sigma} \sum_{\alpha \neq 0} \varepsilon^{-2|\alpha|} \int_{B_{\frac{\varepsilon}{2}}} \left|P^{(\alpha)}\left(D\right) D^{\beta} u\right|^2 dx \leq C \sum_{\alpha \neq 0} \varepsilon^{-2|\alpha|} \int_{B_\varepsilon} \left|P^{(\alpha)}\left(D\right) u\right|^2 dx$$

Proof. Let $\beta \in \mathbb{Z}_+^n \cap \sigma\mathcal{H}$, from (1) of definition 4.1, we have

$$\left|\xi^\beta\right| \leq C^\sigma \left(1 + d\left(\xi\right)\right)^\sigma ,$$

hence $\exists C > 0$, $\forall \varepsilon \in \left]0,1\right[$, $\forall \xi \in \mathbb{R}^n$,

$$\varepsilon^\sigma \left|\xi^\beta\right| \leq C^\sigma d_{\sigma,\varepsilon}\left(\xi\right) \tag{3}$$

Multiplying (4.2) by $\left(2\pi\right)^{-n} \left|\widehat{v}\left(\xi\right)\right|$ and integrating with respect to ξ, we obtain

$$\varepsilon^{2\sigma} \int \left|D^\beta v\right|^2 dx \leq C^2 \left|\left|\left|v\right|\right|\right|_{\sigma,\varepsilon}^2 \tag{4}$$

Let $\varphi \in C_0^\infty\left(B_1\right)$ equals 1 in $B_{\frac{1}{2}}$ and apply the estimate (4.3) to $v = P^{(\alpha)}\left(D\right)\left(\varphi^\varepsilon u\right)$, then

$$\varepsilon^{2\sigma} \int \left|P^{(\alpha)}\left(D\right) D^\beta \left(\varphi^\varepsilon u\right)\right|^2 dx \leq C^2 \left|\left|\left|P^{(\alpha)}\left(D\right)\left(\varphi^\varepsilon u\right)\right|\right|\right|_{\sigma,\varepsilon}^2$$

$$\varepsilon^{2\sigma} \sum_{\alpha\neq0} \varepsilon^{-2|\alpha|} \int \left|P^{(\alpha)}\left(D\right) D^\beta \left(\varphi^\varepsilon u\right)\right|^2 dx \leq C^2 \sum_{\alpha\neq0} \varepsilon^{-2|\alpha|} \left|\left|\left|P^{(\alpha)}\left(D\right)\left(\varphi^\varepsilon u\right)\right|\right|\right|_{\sigma,\varepsilon}^2 ,$$

consequently lemma 4.6 gives

$$\varepsilon^{2\sigma} \sum_{\alpha\neq0} \varepsilon^{-2|\alpha|} \int \left|P^{(\alpha)}\left(D\right) D^\beta \left(\varphi^\varepsilon u\right)\right|^2 dx \leq C \sum_{\alpha\neq0} \varepsilon^{-2|\alpha|} \int_{B_\varepsilon} \left|P^{(\alpha)}\left(D\right)\left(u\right)\right|^2 dx$$

As $\varphi^\varepsilon\left(x\right) = \varphi\left(\frac{x}{\varepsilon}\right) = 1$ in $B_{\frac{\varepsilon}{2}}$, then

$$\varepsilon^{2\sigma} \sum_{\alpha\neq0} \varepsilon^{-2|\alpha|} \int_{B_{\frac{\varepsilon}{2}}} \left|P^{(\alpha)}\left(D\right) D^\beta u\right|^2 dx \leq C \sum_{\alpha\neq0} \varepsilon^{-2|\alpha|} \int_{B_\varepsilon} \left|P^{(\alpha)}\left(D\right)\left(u\right)\right|^2 dx$$

□

Proposition 8. *Let Ω be a bounded open set in $\mathbb{R}^n$ and $\beta \in \mathbb{Z}_+^n \cap \sigma\mathcal{H}$, then there exists a constant $C > 0$, such that for every u solution of $Pu = 0$ in Ω and $\delta \in \left]0,1\right[$, we have*

$$\sum_{\alpha\neq0} \delta^{-2|\alpha|} \int_{\Omega_\delta} \left|P^{(\alpha)}\left(D\right) D^\beta u\right|^2 dx \leq C\delta^{-2\sigma} \sum_{\alpha\neq0} \delta^{-2|\alpha|} \int_{\Omega} \left|P^{(\alpha)}\left(D\right) u\right|^2 dx,$$

where

$$\Omega_\delta = \left\{x \in \Omega : dist\left(x, \partial\Omega\right) > \delta\right\}$$

Proof. The proof is obtained from the precedent lemma and follows the same reasoning as the proof of theorem 4.4.2 of [7]. □

Corollary 9. *Let $P\left(D\right)$ an hypoelliptic operator, then $\exists C > 0$ such that for every solution of $P\left(D\right)u = 0$ in Ω,$\forall \varepsilon \in \left]0,1\right[$, $\forall j = 1, 2, ...$, we have*

$$\varepsilon^{2\sigma} \sum_{0\neq\alpha\in\mathbb{N}_0^n} \varepsilon^{-2|\alpha|} \left\|Q_{\mathcal{H}}\left(D\right) P^{(\alpha)}\left(D\right) u\right\|_{L^2\left(\Omega_{\varepsilon j}\right)}^2 \leq C \sum_{0\neq\alpha\in\mathbb{N}_0^n} \varepsilon^{-2|\alpha|} \left\|P^{(\alpha)}\left(D\right) u\right\|_{L^2\left(\Omega_{\varepsilon(j-1)}\right)}^2 \tag{5}$$

The principal result of this section is the following theorem.

Theorem 10. *Let u be a solution of the hypoelliptic equation $P\left(D\right)u = 0$ in Ω, then for every $\omega \subset\subset \Omega$, there is a constant $C > 0$, such that $\forall j \in \mathbb{N}$, we have*

$$\left\|Q_{\mathcal{H}}^j\left(D\right) u\right\|_{L^2(\omega)} \leq C^{(j+1)} j^{\sigma j} \tag{6}$$

Proof. Since $\rho=\rho(\omega,\partial\Omega)>0$, then there exists $\delta\in\,]0,\rho[$ such that $\omega\subset\Omega_\delta\subset\Omega$. Take $\varepsilon=\dfrac{\delta}{j}$, $j\in\mathbb{N}$, and let us show by induction on j the following estimate

$$\varepsilon^{2j\sigma+2m}\sum_{0\neq\alpha\in\mathbb{N}_0^n}\varepsilon^{-2|\alpha|}\left\|\left(Q_{\mathcal{H}}^{j}(D)\,P^{(\alpha)}(D)\,u\right)\right\|_{L^2(\Omega_{j\varepsilon})}^2<C^{2(j+1)}, \tag{7}$$

where m is the order of P.

As every solution u of an hypoelliptic equation is $\mathcal{C}^\infty$, then there exists $C>0$ such that (4.6) is satisfied for $j=0$. Suppose that (4.6) is true for $j\leq l$ ($l\geq 0$), we have to prove that it remains true for $j=l+1$. Since $v=Q_{\mathcal{H}}^{l}(D)\,u$ is also a solution of equation $P(D)\,u=0$, then from corollary 4.10, we obtain

$$\begin{aligned}&\varepsilon^{2\sigma(l+1)+2m}\sum_{0\neq\alpha}\varepsilon^{-2|\alpha|}\left\|\left(Q_{\mathcal{H}}^{l+1}(D)\,P^{(\alpha)}(D)\,u\right)\right\|_{L^2\left(\Omega_{\varepsilon(l+1)}\right)}^2\\ \leq\;& C\varepsilon^{2\sigma l+2m}\sum_{0\neq\alpha}\varepsilon^{-2|\alpha|}\left\|P^{(\alpha)}(D)\,Q_{\mathcal{H}}^{l}(D)\,u\right\|_{L^2(\Omega_{\varepsilon l})}^2\end{aligned} \tag{8}$$

By the induction hypothesis, we have

$$\varepsilon^{2\sigma l+2m}\sum_{0\neq\alpha}\varepsilon^{-2|\alpha|}\left\|P^{(\alpha)}(D)\,Q_{\mathcal{H}}^{l}(D)\,u\right\|_{L^2(\Omega_{\varepsilon l})}^2\leq C_1^{2(l+1)}\ ,$$

consequently, we obtain

$$\varepsilon^{2\sigma(l+1)+2m}\sum_{0\neq\alpha}\varepsilon^{-2|\alpha|}\left\|\left(Q_{\mathcal{H}}^{l+1}(D)\,P^{(\alpha)}(D)\,u\right)\right\|_{L^2\left(\Omega_{\varepsilon(l+1)}\right)}^2\leq C_2^{2(l+2)},$$

hence $\forall j\in\mathbb{N}$, we have

$$\varepsilon^{2\sigma j+2m}\sum_{0\neq\alpha}\varepsilon^{-2|\alpha|}\left\|P^{(\alpha)}(D)\,Q_{\mathcal{H}}^{j}(D)\,u\right\|_{L^2(\Omega_{\varepsilon j})}^2\leq C_2^{2(j+1)} \tag{9}$$

The estimate (4.8) with $|\alpha|=m$ gives $\forall j\in\mathbb{N}$,

$$\left\|Q_{\mathcal{H}}^{j}(D)\,u\right\|_{L^2(\Omega_{\varepsilon j})}^2\leq\varepsilon^{-2\sigma j}C_2^{2(j+1)},$$

as $\varepsilon=\dfrac{\delta}{j}$, then

$$\left\|Q_{\mathcal{H}}^{j}(D)\,u\right\|_{L^2(\Omega_{\varepsilon j})}^2\leq\left(\frac{j}{\delta}\right)^{2\sigma j}C_2^{2(j+1)}\leq C^{2(j+1)}j^{2\sigma j},$$

hence

$$\left\|Q_{\mathcal{H}}^{j}(D)\,u\right\|_{L^2(\Omega_{j\varepsilon})}\leq C^{(j+1)}j^{\sigma j}$$

□

We denote $G^{s,\,\mathcal{H}}(\Omega)$ the multi-anisotropic Gevrey space associated with $\mathcal{H}$ and by $\mu_{\mathcal{H}}$ and μ_Q the respective formal orders of the Newton's polyhedrons $\mathcal{H}$ and $\Gamma(Q_{\mathcal{H}})$, then we have the following relations

$$\Gamma(Q_{\mathcal{H}})=\sigma\mathcal{H}\ \text{ and }\ \mu_Q=\sigma\mu_{\mathcal{H}}$$

The principal result of this paper is the following theorem.

Theorem 11. *Every solution $u\in\mathcal{D}'(\Omega)$ of the hypoelliptic equation $P(D)\,u=0$ is a function of $G^{\frac{\sigma}{\mu_{\mathcal{H}}},\,\mathcal{H}}(\Omega)$.*

Proof. The theorem 4.11 says that every u solution of the hypoelliptic equation $Pu = 0$ is a Gevrey vector of the operator $Q_{\mathcal{H}}$, i.e. we have $u \in G^{\frac{\sigma}{\mu_Q}}(\Omega, Q_{\mathcal{H}})$. From theorem 3.4 and as the operator $Q_{\mathcal{H}}$ is multi-quasielliptic, then we have $u \in G^{\frac{\sigma}{\mu_Q}, \Gamma(Q_{\mathcal{H}})}(\Omega)$, and consequently $u \in G^{\frac{1}{\mu_{\mathcal{H}}}, \sigma\mathcal{H}}(\Omega)$. A simple computation shows that in general $G^{s, \sigma\mathcal{H}}(\Omega) = G^{s\sigma, \mathcal{H}}\Omega$, hence $u \in G^{\frac{\sigma}{\mu_{\mathcal{H}}}, \mathcal{H}}(\Omega)$. □

Remark 7. *It is interesting to compare the result of the theorem with the microlocal Gevrey regularity result obtained in* [2].

References

[1] C. Bouzar, R. Chaïli, Gevrey vectors of multi-quasielliptic systems, Proc. Amer. Math. Soc. 131 (5) (2003), 1565-1572

[2] C. Bouzar, R. Chaïli, A Gevrey microlocal analysis of multi-anisotropic differential operators. Rend. Sem. Mat. Univ. Pol. Torino, Vol. 64, 3, (2006), 305-318

[3] J. Friberg, Multi-quasielliptic polynomials, Ann. Sc. Norm. Sup. Pisa. Cl. di. Sc. 21 (1967), 239-260.

[4] S. G. Gindikin, L. R. Volevich, The method of Newton polyhedron in the theory of partial differential equations, Kluwer, (1992)

[5] G. H. Hakobyan, Estimates of the higher order derivatives of the solution of hypoelliptic equations. Rend. Sem. Mat. Univ. Pol. Torino, Vol. 61:4, (2003), 443–459.

[6] L. Hörmander, Distributions theory and Fourier analysis, Springer, (1990).

[7] L. Hörmander, Linear partial differential operators, Springer (1969).

[8] V. P. Mikhaïlov, The behavoir at infinity of a class of polynomials, Proc. Steklov. Inst. Mat. 91 (1967), 59-80.

[9] L. Rodino, Linear partial differential operators in Gevrey spaces, World Scientific, (1993).

[10] L. Zanghirati, Iterati di una classe di operatori ipoellipticí e classi generalizzate di Gevrey, Suppl. Boll. U.M.I. (1980), 177-195.

Oran-Essenia University, Algeria
E-mail address: `bouzar@yahoo.com`

University of Bechar, Algeria
E-mail address: `Ahmedalimat@yahoo.fr`